

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/304252604>

Formulation and evaluation of *Abutilon indicum* and *Boerhaavia diffusa* for the determination of nephroprotective activities

Article in *Scientific Transactions in Environment and Technovation* · December 2015

DOI: 10.20894/STET.116.009.002.008

CITATIONS

5

READS

808

1 author:



Velavan Sivanandham

77 PUBLICATIONS 1,006 CITATIONS

SEE PROFILE

Formulation and evaluation of *Abutilon indicum* and *Boerhaavia diffusa* for the determination of nephroprotective activities

S. Shanmugapriya, R.Anuradha, R.Kiruthika, P.Venkatalakshmi and S.Velavan*.

PG and Research Department of Biochemistry, Sengamala Thayaar Educational Trust Women's College, Sundarakkottai, Mannargudi 614 016, Tamil Nadu, India

*Harman Institute of Science Education and Research, Thanjavur, Tamil Nadu, India

Abstract

In vivo anti nephrotoxic activities of the whole plants of *Abutilon indicum* and *Boerhaavia diffusa*, their formulation and individual extracts were determined in *in vivo* models against nephrotoxicity in rats, induced with gentamicin. A control group (water, group I, n = 6) was compared with rats administrated 40 mg/kg gentamicin, once daily for 21 days (Groups II, III, IV, V and VI). The effect of ethanolic extract of *Abutilon indicum* and *Boerhaavia diffusa* formulation and individual extract (Groups IV,V and IV) at a dose level of 200 mg/kg was compared in gentamicin-induced nephrotoxicity. The activities of urea, uric acid, creatinine, sodium and potassium were significantly increased in rats exposed to gentamicin. Administration of ethanolic extract of formulation prevented severe alterations of biochemical parameters. In conclusion, this study obviously demonstrated that pretreatment with ethanolic extract of formulation significantly attenuated the physiological alterations induced by gentamicin. Also, the present study identifies new areas of research for development of better therapeutic agents for kidney diseases.

Keywords: Formulation of *Abutilon indicum* and *Boerhaavia diffusa*, Gentamicin, Nephrotoxicity

INTRODUCTION

Nephrotoxicity is one of the most common kidney problems and occurs when body is exposed to a drug or toxin (Porter and Bennet, 1981). A number of therapeutic agents can adversely affect the kidney resulting in acute renal failure, chronic interstitial nephritis and nephritic syndrome because there is an increasing number of potent therapeutic drugs like aminoglycoside antibiotics, NSAID's, etc., have been added to the therapeutic arsenal in the recent years (Hoitsma, *et al.*, 1991). Exposure to chemical reagents like ethylene, glycol, carbon tetrachloride, sodium oxalate and heavy metals such as lead, mercury, cadmium and arsenic also induces nephrotoxicity. Prompt recognition of the disease and cessation of responsible drugs are usually the only necessary therapy (Paller, 1990). The term renal failure primarily denotes failure of the excretory function of kidney, leading to retention of nitrogenous waste products of metabolism in the blood (Herfindal and Gourley 2000). In addition to this, there is a failure of regulation of fluid and electrolyte balance along with endocrine dysfunction. The renal failure is fundamentally categorized into acute and chronic renal failure (Barry *et al.*, 2000). Acute renal failure (ARF) refers to the sudden and usually reversible loss of renal function which develops over a period of days or weeks. There are many causes for acute renal failure which mainly includes acute tubular necrosis that commonly accounts for 85% of incidence. Mostly acute tubular necrosis occurs either due to ischemia or toxins. The toxins may be exogenous or endogenous. The exogenous agents are radio contrast agents, cyclosporine,

antibiotics, chemotherapeutic agents, organic solvents, acetaminophen and illegal abortifacients (Herfindal and Gourley 2000; Barry *et al.*, 2000).

Chronic renal failure (CRF) is an irreversible deterioration in the renal function which classically develops over a period of years, leading to loss of excretory metabolic and endocrine functions. Various causes of renal failure have been recognized like hypertension, diabetes mellitus, antineoplastic agents like cyclophosphamide, vincristin and cisplatin etc (Herfindal and Gourley 2000).

A number of environmental contaminants, chemicals and drugs including antibiotics dramatically alter the structure and function of various tissues and produce multiple adverse effects in the liver, kidney, heart and intestine (Kohn *et al.*, 2005). Aminoglycoside antibiotics are frequently used in the treatment of severe infections of the abdomen and urinary tract (Nagai and Takano, 2004). Gentamicin (GM) is still considered to be an important aminoglycoside antibiotic against life threatening bacterial infections. However, nephrotoxicity and ototoxicity remain major problems due to its long term clinical use (Khan *et al.*, 2009). GM is known to cause a number of morphologic, metabolic and functional alterations in the kidney and the specificity of GM nephrotoxicity is apparently related to its accumulation in the renal proximal convoluted tubules leading to tubular necrosis (Pedraza-Chaverri, 2000). Aminoglycoside antibiotics have been widely used for gram-negative bacterial infections. However, their nephrotoxicity and ototoxicity are the major limitations in clinical use. Among several aminoglycoside antibiotics, the grade of nephrotoxicity has been reported to be in the following order as, neomycin > gentamicin >

*Corresponding Author :
email:

tobramycin (Hu *et al.*, 1996). Gentamicin nephrotoxicity occurs in about 15-30% of treated subjects, is manifested clinically as non-oliguric renal failure, with a slow rise in serum creatinine and hypoosmolar urinary output developing after several days of treatment (Abdel-zaher *et al.*, 2008). The drug enters into the cells by adsorptive/receptor mediated endocytosis after binding to acidic phospholipids and megalin and is found essentially in lysosomes. Animals treated with low, therapeutically relevant doses of aminoglycosides show both lysosomal phospholipidosis and apoptosis in proximal tubular cells (Suzuki *et al.*, 1995).

Though there are many allopathic drugs have been prescribed to treat nephrotoxicity and related kidney problems, their negative side effects cannot be ignored. Awareness in this regard is growing, and the need of searching for alternative user friendly drugs is often emphasized in various forums and the traditional knowledge and the plants as viable alternative source for finding solution for such ailments have been globally accepted and research is gaining momentum for scientific validation of the traditional knowledge and herbals as the alternative source of therapeutics.(reference).The present article deals with two herbal formulations of *Abutilon indicum* and *Boerhaavia diffusa* with reference to the nephroprotective activities in the gentamicin induced nephrotoxicity in male rats.



Figure 1: *Abutilon indicum*



Figure 2: *Boerhaavia diffusa*

MATERIALS AND METHODS

Plant Collection and Identification

Abutilon indicum (Fig 1) and *Boerhaavia diffusa* (Fig 2) were collected in and around Mannargudi, Thiruvavur (Dt), Tamil Nadu.

Preparation of plant powder

The plants were air dried under shade for 10-15 days. Then the dried materials were grinded to fine powder using an electric grinder and stored in air tight bottles. The powders were used for further determination of nephroprotective activities.

Extraction of plant material

Ethanolic extracts were prepared according to the methodology of Indian pharmacopoeia (Anonymous, 1996). The coarse powder material was subjected to soxhlet extraction separately and successively with ethanol and distilled water. These extracts were concentrated to dryness in flash evaporator under reduced pressure and controlled temperature (40°C-50°C). The ethanolic and aqueous extracts were put in air tight container and stored in refrigerator.

Anti nephrotoxic activity in animal model

Male albino rats of Wistar strain approximately weighing 150-180g were used in this study. They were healthy animals purchased from the Indian Institute of Science, Bangalore. The animals were housed in spacious polypropylene cages bedded with rice husk. The animal room was well ventilated and maintained under standard experimental conditions (Temperature $27 \pm 2^\circ$ C and 12 hour light/dark cycle) throughout the experimental period. All the animals were fed with standard pellet diet and water was provided *ad libitum*. They were acclimatized to the environment for one week prior to experimental use. The animal feed composition included crude protein (22.3%), crude oil (4.01%), crude fibre (4.02%), Ash (8.02%), calcium (1-2%), phosphorous (0-6%) and sand silical (1.02%).

Experimental design

In this experiment, a total of 24 rats were used. The rats were randomly divided into four groups of six rats in each group.

Group I: Normal animal received standard feed and water

Group II: Animals received oral administration of gentamicin (40 mg/kg body weight) for 21 days (Annie *et al.*, 2005)

Group III: Treatment group received gentamicin as group II treated with extract of *Abutilon indicum* at a dose of 200 mg/kg body weight for 21 days.

Group IV: Treatment group received gentamicin as group II treated with furosemide at a dose of 13 mg/kg bodyweight for 21 days (Viviane Gomes Portella *et al.*, 2012).

Group V: Treatment group received gentamicin as group II treated with extract of *Abutilon indicum* at a dose of 200 mg/kg body weight for 21 days.

Group VI: Treatment group received gentamicin as group II treated with formulation (equal concentrations of *Abutilon indicum* and *Boerhaavia diffusa*) at a dose of 200mg/kg body weight for two weeks.

During the experimental period, food and water consumption were measured every day and the body weight was measured at the initial (day 1) and final day (day 21) of the experiment.

Collection of blood and preparation of serum sample

At the end of the experimental period, the animals were anaesthetized using chloroform vapour prior to dissection. Blood was collected by cardiac puncture into serum separator tubes. The blood was allowed to clot by standing at room temperature for 30 minutes and then refrigerated for another 30 minute. The resultant clear part was centrifuged at 3000 rpm for 10 minutes, and then the serum (supernatant) was isolated and stored in refrigerator until further analysis.

RESULTS AND DISCUSSION

In the recent years, although technology and medicine have developed extensively, some countries have made it obligatory to use natural products for treatment diseases and many different purposes. Many other countries have taken initiative for the exploitation of the traditional knowledge and the plants known to people with health benefits and the idea is extensively used for the treatment of various diseases. Indian system of medicinal practices has traditionally used the locally available herbals for the treatment of various ailments. *A. indicum* and *B. diffusa* are the two important herbals with proven medicinal properties. *A. indicum* is reported to have analgesic (Kalyani, 1985;

Ahmed *et al.*, 2000), hepatoprotective (Roshan *et al.*, 2004), hypoglycemic activity (Seetharam *et al.*, 2000), wound healing activity (Roshan *et al.*, 2008), antidiabetic (Adisakwattana *et al.*, 2009), and anti diarrheal (Chandrashekar *et al.*, 2004). Different parts of the *B. diffusa* have been widely used for the treatment of dyspepsia, jaundice, enlargement of spleen, abdominal pain, abdominal tumors, and urinary disorders used in the traditional medicine (Kersten *et al.*, 1998). Pharmacological studies have demonstrated that *B. diffusa* known to possess diuretic (Kirtikar and Basu, 1956); nephrotic syndrome (Gaitonde and Kulkarn, 1974); anti-inflammatory and anti-nociceptive (Singh and Udupa, 1972); anticonvulsant (Hiruma-Lima *et al.*, 2000); immunomodulatory (Kaur and Goel, 2009); hepatoprotective (Mungantiwar *et al.*, 2001); antiurolithiatic (Rawat and Mehrotra, 1997); antioxidant and antidiabetic activity (Pareta and Patra, 2011). The whole plant analysis of *B. diffusa* is known to contain numerous phytochemical constituents that include flavonoids, alkaloids, triterpenoids, steroids, lipids, lignins, tannins, phlobaphenes and ursolic acid (Satheesh and Pari, 2004; Jain and Singh, 1994). The present investigation reveals the following findings *In vivo* anti nephrotoxic activity

Nephroprotective activities of *Abutilon indicum* and *Boerhaavia diffusa* formulation and individual extract were as follows

Changes in renal functional markers

The effects of the plants on the selected kidney functions are presented in Table 1. It represents the levels of uric acid, potassium, sodium, urea and creatinine in serum of experimental rats. Group II gentamicin intoxicated rats showed a significant increase in the level of uric acid, potassium, sodium, urea and creatinine when compared to Group I rats. Group III gentamicin intoxicated rats treated with *Abutilon indicum* extract and Group IV gentamicin intoxicated rats treated with furosemide showed significant decrease in the level of uric acid, potassium, sodium, urea and creatinine when compared to group II.

Table 1: Effect of extract of *Abutilon indicum* and *Boerhaavia diffusa* formulation and individual extract on gentamicin induced nephrotoxicity in experimental rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
Creatinine (mg/dl)	0.95±0.232#	2.65±0.328*	0.955±0.152#	1.38±0.177 #	0.95±0.15#	1.21±0.215 #
Urea (mg/dl)	46.66±13.23#	140.94±17.58*	49.52±12.34#	81.9±9.33 #	40.9±3.98#	65.71±10.69#
Uric Acid (mg/dl)	2.23±0.891 #	7.62±1.251 *	2.88±0.71 #	4.226±0.91 #	4.23±1.07#	4.035±0.721#
Potassium (Meq/L)	4.03±0.248 #	9.28±0.12 *	5.09±0.02 #	6.89±0.18 #	5.43±0.15#	5.89±0.17#
Sodium (Meq/L)	151.69±3.48#	175.76±4.32 *	148.60±1.73#	155.54±2.41#	143.3±2.25#	152.46±1.99#

*Significantly different from Group I, III and IV (p<0.05); # Significantly different from Group II (p<0.05)

Urea is an end product of protein catabolism. It is freely filtered by the glomerulus, passively reabsorbed in both the proximal and distal nephron and excreted in high concentration in urine. The excretion of urea was recognized as an estimate of kidney function. The serum urea level is used as an index of kidney function (Lesley and Lvey, 2005). Drugs that can increase urea levels include allopurinol, some diuretics, gentamicin and indometacin (Mason, 2004). In the present study also observed that the increased level of urea in gentamicin intoxicated rats. Supplementation of formulation restored the increased level of urea in gentamicin-induced rats. This result is in line with the findings of Ali Noorani *et al.* (2011) and Kotnis *et al.* (2004).

Creatinine is an end product of muscle catabolism, which is removed at a constant rate by the kidneys. The concentration of creatinine in serum is the most widely used and commonly accepted measure of renal function in clinical medicine. The clinical utility of the serum creatinine concentration centers on its relation to the glomerular filtration rate (GFR) (Perrone *et al.*, 1992). The serum creatinine concentration is the most commonly used index of the kidney function. The level of creatinine in the blood rises if the kidney does not function properly (Lesley and Levey, 2005). Gentamicins have been reported to increase creatinine measurements (Mason, 2004). In the present study also it was found that the creatinin level was increased in response to gentamycin. Administration of formulation restored the level of creatinine in gentamicine treated rats. This study is consisted with Sara *et al.* (2009) and Kotnis *et al.* (2004) studies

Uric acid is the end product of purine degradation. It is produced by xanthine oxidase from xanthine which in turn is produced from purine. It is sparingly soluble in water. Degradation of uric acid mainly takes place in liver. Elevated serum uric acid is correlated inversely with renal blood flow/m²body surface area and directly with renal vascular and total resistance and metabolic syndrome with or without a low globular filtration rate (Suchetha Kumari *et al.*, 2011).

Sodium is the most abundant cation in the extracellular fluid and is the major regulating factor for bodily water balance. Extracellular (i.e., intravascular and interstitial) and intracellular sodium contents are closely affected by the body fluid status. The kidneys are the primary organ responsible for the retention and excretion of body sodium and water. Potassium is the primary cation in the intracellular space, with an average intracellular fluid concentration of about 140 mEq/L (140 mmol/L) (Rose, 2001). The major physiological role of potassium is in the regulation of muscle and nerve excitability. It may also play important roles in the control of intracellular volume (similar to the ability of sodium in controlling extracellular volume), protein synthesis,

enzymatic reactions and carbohydrate metabolism (Zull, 1989). Acid, bases and salts are collectively called electrolytes. Electrolyte imbalance can lead to serious consequences as it affects the homeostasis of the body. Homeostasis is the process by which the body cells maintain their internal balance in spite of changes in the external environment, commonly measured electrolytes are sodium, potassium, calcium, chloride, bicarbonate, etc., which are good indicators of kidneys function (Cohen and Lemann, 1991). In the present study, gentamicin treated rats showed significantly increased sodium and potassium levels when compared to normal control rats. Administration of formulation restored the normal level of sodium and potassium in gentamicin treated rats. This result coincides with Ali Noorani *et al.* (2011).

The ethanolic extract at equal concentrations of formulation showed maximum nephroprotective activity compared to furosemide. On the basis of the results obtained in the present study, it is concluded that the formulation has potent nephroprotective activity.

ACKNOWLEDGEMENTS

The authors are grateful to the Management of Sengamala Thayaar Educational Trust Women s College, Sundarakkottai, Mannargudi for the facilities and encouragement and support.

REFERENCES

- Abdel-Zaher AO, Abdel-Hady RA, Mahmoud MM, Farrag MMY. The potential protective role of alphasilicic acid against acetaminophen- induced hepatic and renal damage. *Toxicol.*, 2008; 243: 261-270.
- Adisakwattana S, Pudhom K, Yibcho,S.2009. Influence of the methanolic extract from *Abutilon indicum* leaves in normal and streptozotocin-induced diabetic rats. *African J of Biotechnology*, ; 8(10): 2011-2015.
- Ahmed M, Amin S, Islam M, Takahashi M, Okuyama E. Analgesic principal from *Abutilon indicum*. *Pharmazie*, 2000;55(4): 314-316.
- Al-Shabanah OA, Aleisa AM, Al-Yahya AA, Al-Rejaie SS, Bakheet SA, Fatani AG, et al. Increased urinary losses of carnitine and decreased intramitochondrial coenzyme A in gentamicin-induced acute renal failure in rats. *Nephrol Dial Transplant*, 2010; 25:69-76.
- Annie, S., Rajagopal, P.L., Malini, S., 2005. Effect of *Cassia auriculata* Linn. Root extract on cisplatin and gentamicin-induced renal injury. *Phytomedicine*, 12, 555-560.
- Annie, S., Rajagopal, P.L., Malini, S., 2005. Effect of *Cassia auriculata* Linn. Root extract on cisplatin and gentamicin-induced renal injury. *Phytomedicine*, 12, 555-560.

- Anonymous. The Wealth of India, Raw Materials Vol. 2B, Publication and Information Directorate, CSIR New Delhi: 1996; 90.
- Arshad Ali Noorani, Karan Ajay Gupta, Khushboo Bhadada and M. K. Kale. Protective Effect of Methanolic Leaf Extract of *Caesalpinia Bonduc* (L.) on Gentamicin-Induced Hepatotoxicity and Nephrotoxicity in Rats. Iranian Journal of Pharmacology and Therapeutics. 2011; 10 (1) : 21-25.
- Barry M, Brenner, Floyd C, Rector. The kidney 6th Ed. Vol I, W.B. Saunders Company, Philadelphia; 2000; 3-67.
- Boneses RN and Tausk HA. On the colorimetric determination of creatinine by the Jaffe reaction. *J. Biol. Chem.* 1945; 158: 581-591.
- Chandrashekhar VM, Nagappa AN, Channesh TS, Habbu PV, Rao KP. Anti-diarrhoeal activity of Abutilon indicum Linn leaf extracts. *J of Natural Remedies*, 2000;4: 12-16.
- Cohen EP and Lemann J. The role of the laboratory in evaluation of kidney function. *Clinical chemistry*. 1991; 37 (6): 785-796.
- del Valle DAS, Imbrogno MA, Fernadez E. Gentamicin in pediatric infections Hendriks JGE, van Horn JR, van der Mei HC, Busscher HJ. Backgrounds of antibiotic loaded bone cement and prosthesis-related infection. *Biomaterials*, 2004; 25: 545-56.
- Fossati P Prencipe L and Berti G. Use of 3,5-dichloro-2-hydroxy benzene sulfonic and/4-aminophenazone chromogenic system in direct enzymic assay of uric acid in serum and urine. *Clin chem.*, 1980; 26: 227-231.
- Gaitonde BB, H. Kulkarn, Nabar SD. Diuretic activity of punarnava (Boerhaavia diffusa). *Bull. Haffkine Inst.*, 1974; 2: 24-25.
- Herfindal, Gourley. Text book of therapeutic drug and disease management. 7th Edn. Charcil Livingstone, London; 2000; 425-36.
- Hiruma-Lima CA, Gracioso JS, Bighetti EJ. The juice of fresh leaves of Boerhaavia diffusa L. (Nyctaginaceae) markedly reduces pain in mice. *J. Ethnopharmacol.*, 2000; 71: 267-274.
- Hoitsma AJ, Wetzels JF and Koene RA. Drug induced nephrotoxicity. Aetiology, clinical features and management, *Drug Saf.*, 1991; 6 (2): 131-147.
- Hu J J, J.S.H. Yoo, M. Lin, E-J Wang, C.S Yang. Protective effects of diallyl disulfide on Acetaminophen induced toxicities. *Food Chem Toxicol.*, 1996; 34: 963-969.
- Jain S.P. and Singh S.C., Ethnomedical botanical survey of Ambikapur District, M.P. Ethnobiology in human welfare. Abstracts of the 4th International Congress of Ethnobiology, Lucknow, Uttar Pradesh, India, 1994; 293.
- Kalyani GA. A preliminary pharmacological of Abutilon indicum analgesic activity. *Fitoterapia*, 1985;56 (3): 169-171.
- Kaur M, Goel RK. Anti-convulsant Activity of Boerhaavia diffusa: Plausible Role of Calcium Channel Antagonism, *Evid. Comp. Alt. Med.* 2009; 1-7.
- Kersten L, Braunlich H, Kepper BK, Gliesing C, Wendelin M, Westphal J. *J appltoxicol.* 1998; 8(2): 245
- Khan SA, Priyamvada S, Farooq N, Khan S, Wasim KM, Yusufi ANK. Protective effect of green tea extract on gentamicin-induced nephrotoxicity and oxidative damage in rat kidney. *Pharmacol Res.*, 2009; 59: 254-262.
- Kirtikar KR, Basu BD. Indian Medicinal Plants Vol. III, Lalit Mohan Basu, Allahabad, 1956, 2045-2048.
- Kirtikar KR, Basu BD. Indian Medicinal Plants, Edn 2, Vol. I, Dehradun 1994, 314-317.
- Kohn S, Fradis M, Robinson E, Iancu TC. Hepatotoxicity of combined treatment with cisplatin and gentamicin in the guinea pig. *Ultrastruct Pathol.*, 2005; 29: 129-137.
- Kotnis MS, Patel P, Menon SN and Sane RT. (2004) Renoprotective effect of Hemidesmus indicus, a herbal drug used in gentamicin induced renal toxicity. *Nephrology (Carlton)*. 9(3); 142-152.
- Lesley AS and Levey AS. (2005) Measurements of Kidney function. *Medical Clinical North America*. 89: 457-473.
- Martinez-Salgado C, Lopez-Hernandez FJ, Lopez-Novoa JM. Glomerular nephrotoxicity of aminoglycosides. *Toxicol Appl Pharmacol.*, 2007; 223: 86-98.
- Maruna RFL. (1957) Determination of serum potassium by colorimetric method. *Clinica Chemica Acta*, 2(2): 131-133.
- Maruna RF and Trinder SR (1958) Determination of serum sodium by colorimetric method. *Clin. Chem Acta*, 2.1.581.
- Mungantiwar A, Nair AM, Shinde A, Dikshit AJ. Studies on the immunomodulatory effects of Boerhaavia diffusa alkaloidal fraction. *J. Ethnopharmacol.*, 2001; 65: 125-131.
- Nadkarni AK. Indian Materia Medica, Popular Prakashan (Pvt) Ltd, Bombay, 1995, 8-9.
- Nagai J, Takano M. Molecular aspects of renal handling of aminoglycosides and strategies for preventing the nephrotoxicity. *Drug Metab Pharmacokin.*, 2004; 19: 159-170.
- Natelson S. (1957) Micro-techniques of clinical chemistry for the routine laboratory. C.C.Thomas, Springfield, Illinois, 381.
- Paller MS., Drug induced nephropathies. *Med Clin North Am*, 1990; 74 (4): 909-917.
- Pareta SK, Patra KC, Mazumder PM, Sasmal D. Boerhaavia diffusa linn aqueous extract as curative agent in ethylene glycol induced urolithiasis. *Pharmacologyonline.*, 2011, 3: 112-120.

- Pedraza-Chaverri J, Maldonado PD, Medina-Campos ON, Olivares-Corichi IM, Grandos-Silvestre M, Hernandez-Pando R, et al. Garlic ameliorates gentamicin nephrotoxicity: relation to antioxidants. *Free Radic Biol Med.*, 2000; 29: 602-611.
- Porter GA and Bennett WM. Nephrotoxic acute renal failure due to common drugs. *American journal of Physiology*, 1981; 241(7): F1-F8.
- Prajapati ND, Purohit SS, Sharma AK, Kumar TA. Handbook of Medicinal Plants, AGROBIOS, (India) Jodhpur, 2003, 3.
- Rawat AK, Mehrotra S. Hepatoprotective activity of *Boerhaavia diffusa* L. roots: a popular Indian ethnomedicine. *J. Ethnopharmacol.*, 1997; 56: 61-66.
- Rose BD. Clinical Physiology of Acid-Base and Electrolyte Disorders. 5th ed. New York, NY: McGraw-Hill; 2001.
- Roshan S, Ali Sadath, Saleem MA, Patel MJ, Ali Asgar. Effect of *Abutilon indicum* on carbon tetrachloride induced hepatic damage in albino rats. *J Pharma Res.*, 2004; 3(4): 61-62.
- Roshan S, Sadath Ali, Abdullah Khan, Tazneem B, Purohit MG. Wound Healing activity of *Abutilon indicum*. *Phcog Mag* 2008; 4(15): 85-88.
- Sandhu JS, Sehgal A, Gupta O, Singh A. Aminoglycoside nephrotoxicity revisited. *J Ind Acad Clin Med.*, 2007; 8: 331-3.
- Sara A. Khan, Shubha Priyamvada, Neelam Farooq, Sheeba Khan, Md Wasim Khan, Ahad N.K. Yusuf. Protective effect of green tea extract on gentamicin-induced nephrotoxicity and oxidative damage in rat kidney. *Pharmacological Research*, 2009; 59 : 254-262
- Satheesh MA, Pari L. Antioxidant effect of *Boerhavia diffusa* L. in tissues of alloxan induced diabetic rats. *Indian J. Exp. Biol.*, 2004, 42: 989-92.
- Schrier RW, Gottschalk CW. Disease of kidney, 5thed, 2, Published by Little Brown & Co, 1993; 1031-1165.
- Seetharam G, Halageri, Setty SR, Bheemachar. Hypoglycemic activity of *Abutilon indicum* leaf extract in rats. *Fitoterapia*, 2000; 73(2): 156-159.
- Silan C, Uzun O, Ustundag N, Gookcen S, Bedirhan S and Cengiz M. Gentamicin induced nephrotoxicity in rats ameliorated and healing effects of resveratrol. *Biol. Pharm. Bull.*, 2007; 30(1): 79-83.
- Singh RH, Udupa KN. Studies on the Indian indigenous drug punarnava (*Boerhaavia diffusa* Linn.). Part IV: Preliminary controlled clinical trial in nephrotic syndrome. *J. Res. Indian Med.*, 1972; 7: 28-33.
- Suchetha Kumari N, Atul Kamath M, Sukanya Shetty, Roopa Rani Bhandary, Kathyayini. Serum uric acid as obesity related indicator in young obese adults. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2011; 2(2): 1-5.
- Suzuki, S.S., Takamura, J., Yoshida, Y., Shinzawa, Niwat, Tamatani, S R., Comparison of gentamicin nephrotoxicity between rats and mice. *Camp. Biochem. Physiol.*, 1995; 1: 15-28.
- Viviane Gomes Portella, Gustavo Pereira Cosenza, Lucio Ricardo Leite Diniz, Lillian Fernanda Pacheco, Giovanni Dantas Cassali, Marcelo Vidigal Calari, Maria das Graças Lins Brandao, Maria Aparecida Ribeiro Vieira. Nephroprotective Effect of *Echinodorus macrophyllus* Micheli on Gentamicin-Induced Nephrotoxicity in Rats. *Nephron Extra*. 2012; 2: 177-183.
- Zull DN. Disorders of potassium metabolism. *Emerg Med Clin North Am*. 1989; 7: 771-94.